gui_ge

10ug

biao_da_su_zhu

E.coli

biao_qian

NA

fen_zi_liang

6.2 KDa

biao_guan_fen_zi_liang

11 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of 20mM Tris, 200mM NaCl, pH 8.0.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.001 ng/μg (0.01 EU/μg) as determined by LAL test.

huo_xing

Measured in a cell proliferation assay using BALB/c 3T3 cells. The ED50 for this effect is 60-450 pg/ml.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Epidermal growth factor (EGF) is a small 53 amino acid residue long protein that contains three disulfide bridges. It is a small mitogenic protein that is thought to be involved in mechanisms such as normal cell growth, oncogenesis, and wound healing. EGF stimulates the growth of various epidermal and epithelial tissues in vivo and in vitro and of some fibroblasts in cell culture. This protein shows both strong sequential and functional homology with human type-alpha transforming growth factor (hTGF alpha), which is a competitor for EGF receptor sites.

本产品仅供科研使用，不用于临床诊断。