Recombinant Mouse IL-2

货号：EPT026

gui_ge

10ug

biao_da_su_zhu

E.coli

biao_qian

NA

fen_zi_liang

17.4 KDa

biao_guan_fen_zi_liang

17 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of 20mM Sodium Citrate, 0.2% Tween 80, pH 3.0.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.001 ng/μg (0.01 EU/μg) as determined by LAL test.

huo_xing

Loaded Human IL-2RA-Fc(Cat#CJ78) on Protein A Biosensor, can bind Mouse IL-2(Cat#CK24) with an affinity constant of 0.17 uM as determined in BLI assay.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Interleukin 2 (IL 2), also termed T-cell growth factor, is a member of the cytokine family which includes IL-4, IL-7, IL-9, IL-15 and IL-21. Each member of this family has a four alpha helix bundle. IL-2 signals through the IL-2 receptor, a complex consisting of tree subunits, termed alpha, beta and gamma. The IL-2 R gamma is

shared by cytokine receptors of all members of cytokine family. Mature mouse IL2 shares 56% and 73% aa sequence identity with human and rat IL2, respectively. IL-2 is produced by CD4+ T cell, CD8+ T cells, gamma δ T cells, B cells, dendritic cells and eosinophils, and plays a vital role in key function of the immune system, tolerance and immunity, primarily via its potent stimulatory activity for T cells.Thus, IL2 may be a key cytokine in the natural suppression of autoimmunity.

本产品仅供科研使用，不用于临床诊断。