Recombinant Human CD207 (N-6His)

货号: EPT060

gui_ge

50ug

biao_da_su_zhu

Human Cells

biao_qian

N-6His

fen_zi_liang

31.5 KDa

biao_guan_fen_zi_liang

30-40 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Langerin (CD207) is a type II transmembrane glycoprotein which is member K of the C-type lectin domain family. Langerin is used as a marker for Langerhans cells (LCs) which represent the immature dendritic cells in the epidermis. Langerin is necessary and sufficient for Birbeck granule formation. Human langerin sequence contains a 43 aa cytoplasmic domain, a 21 aa transmembrane domain and a 264 aa extracellular domain (ECD) that contains a coiled-coil domain and a single C-type lectin domain. Human langerin shares 68%, 62%, 71% aa identity with mouse, rat and bovine langerin ECD, respectively. Trimerization greatly increases the lectin binding affinity. Langerin internalizes endogenous proteins such as type I procollagen. Internalization by LC is thought to lead to suppression of self reactions. Langerin also mediates endocytosis of non-peptide antigens containing mannose, N-acetyl glucosamine and fucose that are expressed by mycobacteria and fungae.

本产品仅供科研使用，不用于临床诊断。