Recombinant 2019-nCoV S1 Protein

货号: RB91EPT025

gui_ge

50ug

biao_da_su_zhu

Human Cells

biao_qian

NA

fen_zi_liang

78.3kDa

biao_guan_fen_zi_liang

100-130kDa, reducing conditions

xing_tai

Supplied as a 0.2 μm filtered solution of PBS, pH 7.4

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

huo_xing

Immobilized 2019-nCoV S1 Protein(Cat#DRA35) at 2μg/ml (100 μl/well) can bind Human

ACE-2-Fc(Cat#C05Y). The ED50 of Human ACE-2-Fc(Cat#C05Y) is 15-150 ng/ml.

chu_cun_fang_fa

Reconstituted protein solution should be stored at ≤ -20° C.

yun_shu_tiao_jian

The product is shipped on dry ice pack.Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. Most notable is severe acute respiratory syndrome (SARS). The severe acute respiratory syndrome-coronavirus (SARS-CoV) spike (S) glycoprotein alone can mediate the membrane fusion required for virus entry and cell fusion. It is also a major immunogen and a target for entry inhibitors. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion.The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

本产品仅供科研使用，不用于临床诊断。