Recombinant Rat VEGF 164

货号: RB91EPT063

gui_ge

10ug

biao_da_su_zhu

P.Pichia

biao_qian

NA

fen_zi_liang

19.2 KDa

biao_guan_fen_zi_liang

18-23 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

huo_xing

Immobilized Rat VEGF 164(Cat#CJ96) at 2μg/ml (100 μl/well) can bind Mouse VEGFR2-Fc(Cat#C01A).The ED50 of Mouse VEGFR2-Fc(Cat#C01A) is 22.17 ng/ml.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Vascular endothelial growth factor (VEGF/VEGF-A) is originally known as vascular permeability factor (VPF). It belongs to the PDGF family with a cysteine-knot structure comprised of eight conserved cysteine residues, and reckoned as a potent mediator in the process of angiogenesis and vasculogenesis in either fetus or adult. VEGF is particularly expressed in supraoptic , paraventricular nuclei and the choroid plexus of the pituitary, and abundant in the corpus luteum of the ovary and in kidney glomeruli. The rat VEGF protein contains a putative 20 amino acids (aa) signal peptide, and alternative splicing of rat VEGF gene produces isoforms of 120, 144, 164 and 188 aa. Rat VEGF164 respectively displays 97% and 88% aa identity with that regions of mouse and human VEGF. VEGF can bind to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin, and play important roles in inducing endothelial cell proliferation, promoting cell migration, inhibiting apoptosis and inducing permeabilization of blood vessels.

本产品仅供科研使用，不用于临床诊断。