Recombinant Human/Mouse/Rat Irisin (C-Fc)

货号: RB91EPT075

gui_ge

10ug

biao_da_su_zhu

Human Cells

biao_qian

C-Fc

fen_zi_liang

39.7 KDa

biao_guan_fen_zi_liang

50-60 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Fibronectin type III domain-containing protein 5, the precursor of irisin, is a protein that is encoded by the FNDC5 gene.Human Irisin is synthesized as a 212 amino acid (aa) precursor encoding a type 1 transmembrane protein with a 121 aa extracellular domain (ECD), a 21 aa transmembrane domain, and a 39 aa cytoplasmic domain. The ECD of Irisin contains a fibronectin type III domain and multiple glycosylation sites. The ECD is proteolytically cleaved to release the 112 aa soluble Irisin hormone into circulation.Mature human, mouse share 100% sequence identity.Irisin induces expression of peroxisome proliferatoractivated receptor γ coactivator 1α (PGC1α) and uncoupling protein1(UCP1), mitochondrialassociated metabolic proteins. Irisin induces the transition of white adipose tissue into more metabolically active beige adipose tissue.Irisin also regulates neuronal cell differentiation and neurite outgrowth in the brain and is involved in the differentiation of osteoblasts.

本产品仅供科研使用，不用于临床诊断。