Recombinant Cynomolgus TSLP (C-6His)

货号: RB91EPT093

gui_ge

10ug

biao_da_su_zhu

E.coli

biao_qian

C-6His

fen_zi_liang

16.2 KDa

biao_guan_fen_zi_liang

16 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of 20mM PB, 500mM NaCl, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

huo_xing

Immobilized Anti-Human TSLP mAb (Cat#NC066) at 2μg/ml (100 μl/well) can bind Cynomolgus TSLP-His (Cat#CR62). The ED50 of Cynomolgus TSLP-His (Cat#CR62) is 4.48 ng/ml.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Thymic stromal lymphopoietin (TSLP) is a protein belonging to the cytokine family, contains 140 amino acids. It is known to play an important role in the maturation of T cell populations through activation of antigen presenting cells. TSLP induces the release of T-cell-attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c+ dendritic cells. It can induce allergic inflammation by directly activating mast cells. TSLP is produced mainly by non-hematopoietic cells such as fibroblasts, epithelial cells and different types of stromal or stromal-like cells. These cells are located in regions where TSLP activity is required.

本产品仅供科研使用，不用于临床诊断。