Recombinant Human NPTX1 (C-6His)

货号: RB91EPT103

gui_ge

10ug

biao_da_su_zhu

Human Cells

biao_qian

C-6His

fen_zi_liang

45.9 KDa

biao_guan_fen_zi_liang

50-55 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of PBS, 1mM EDTA, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Neuronal Pentraxin (NPTX1, NP1) is a secreted glycoprotein within the Pentraxin family. NPTX1 is co-expressed and forms heteromultimers with the related secreted protein, NPTX2/NARP, NPTXR (Neuronal Pentraxin Receptor) at excitatory synapses. Mature human NPTX1 shares 97% aa sequence identity with mouse, and rat NPTX1. It is produced by hippocampal, cerebral and cerebellar neurons, retinal ganglia and the inner nuclear layer of the retina. It is enriched on presynaptic axonal membranes where it forms complexes with NPTXR. It may be involved in mediating uptake of synaptic material during synapse remodeling or in mediating the synaptic clustering of AMPA glutamate receptors at a subset of excitatory synapses.

本产品仅供科研使用，不用于临床诊断。