Recombinant Mouse LR3-IGF-1 (C-6His)

货号: RB91EPT112

gui_ge

50ug

biao_da_su_zhu

E.coli

biao_qian

C-6His

fen_zi_liang

10.2 KDa

biao_guan_fen_zi_liang

12 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of 20mM NaAc, pH 4.5.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Insulin-like growth factor I (IGF1) belongs to the family of insulin-like growth factors that are structurally homologous to proinsulin. Mouse IGF-I is synthesized as two precursor isoforms with alternate N- and C-terminal propeptides. These isoforms are differentially expressed by various tissues. Mature mouse IGF-I shares 94% and 99% aa sequence identity with human and rat IGF-I, respectively, and exhibits cross-species activity. It shares 60% aa sequence identity with mature mouse IGF-II. IGF-I induces the proliferation, migration, and differentiation of a wide variety of cell types during development and postnatally. It plays an important role in muscle regeneration and tumor progression. IGF-I binds IGF-I R, IGF-II R, and the insulin receptor. IGF-I association with IGF binding proteins increases its plasma half-life and modulates its interactions with receptors.

本产品仅供科研使用，不用于临床诊断。