Recombinant Mouse CD47 (C-Fc)

货号: RB91EPT116

gui_ge

10ug

biao_da_su_zhu

Human Cells

biao_qian

C-Fc

fen_zi_liang

42.8 KDa

biao_guan_fen_zi_liang

60-90 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

CD47, also known as Integrin-Associated Protein (IAP) and OA3, is a glycosylated atypical member of the immunoglobulin superfamily. Mouse CD47 is an integral membrane protein that consists of a extracellular domain (ECD) with a single Ig-like domain, five membrane-spanning regions with short intervening loops, and C-terminal cytoplasmic tail. CD47 has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins. It plays an important role in memory formation and synaptic plasticity in the hippocampus. As a receptor for SIRPA, it binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. Interaction with SIRPG mediates cellcell adhesion, it enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation. It may play a role in membrane transport and/or integrin dependent signal transduction. It also prevents premature elimination of red blood cells.

本产品仅供科研使用，不用于临床诊断。