gui_ge

50ug

biao_da_su_zhu

E.coli

biao_qian

NA

fen_zi_liang

9 KDa

biao_guan_fen_zi_liang

14 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of 10mM HAc-NaAc, 150mM NaCl, 5% Mannitol, pH 4.0.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Parathyroid hormone is the most important endocrine regulator of calcium and phosphorus concentration in extracellular fluid. This hormone is secreted from cells of the parathyroid glands and finds its major target cells in bone and kidney. Another hormone, parathyroid hormone-related protein, binds to the same receptor as parathyroid hormone and has major effects on development. Like most other protein hormones, parathyroid hormone is synthesized as a preprohormone. After intracellular processing, the mature hormone is packaged within the Golgi into secretory vesicles, the secreted into blood by exocytosis. Parathyroid hormone is secreted as a linear protein of 84 amino acids.

本产品仅供科研使用，不用于临床诊断。