Recombinant Human IL-2 (aldesleukin)

货号: RB91EPT147

gui_ge

10ug

biao_da_su_zhu

E.coli

biao_qian

aldesleukin

fen_zi_liang

15.5 KDa

biao_guan_fen_zi_liang

14 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of 10mM Acetata-Na, 5% Trehaiose, pH 4.5.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

huo_xing

Measured in a cell proliferation assay using CTLL-2 mouse cytotoxic T cells. The ED50 for this effect is 90-270 pg/ml.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Recombinant Human Interleukin-2 is a highly purified protein with a molecular weight of approximately 15,300 Daltons. The chemical name is des-alanyl-1, serine-145 Human Interleukin-2. It is produced by recombinant DNA technology using a genetically engineered E. coli strain containing an analog of the human interleukin-2 gene. Genetic engineering techniques were used to modify the Human IL-2 gene, and the resulting expression clone encodes a modified Human IL-2. This recombinant form differs from native Interleukin-2 in following ways: it is not glycosylated; the molecule has serine substituted for cysteine at amino acid position 145; the aggregation state of molecule is likely to be different from that of native IL-2.

本产品仅供科研使用，不用于临床诊断。