Recombinant Human PRDX1 (N, C-6His)

货号: RB91EPT198

gui_ge

50ug

biao_da_su_zhu

E.coli

biao_qian

N, C-6His

fen_zi_liang

25.3 KDa

biao_guan_fen_zi_liang

26 KDa, reducing conditions

xing_tai

Supplied as a 0.2 μm filtered solution of PBS, 10% Glycerol, 0.1mM DTT, pH 6.0.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

chu_cun_fang_fa

Store at ≤-70° C, stable for 6 months after receipt.
Store at ≤-70° C, stable for 3 months under sterile conditions after opening.
Please minimize freeze-thaw cycles.

yun_shu_tiao_jian

The product is shipped on dry ice/polar packs.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Peroxiredoxin-1(PRDX1) contains 1 thioredoxin domain and belongs to the AhpC/TSA family. PRDX1 constitutively expressed in most human cells and it is induced to higher levels upon serum stimulation in untransformed and transformed cells. PRDX1 is involved in redox regulation of the cell. It reduces peroxides with reducing equivalents provided through the thioredoxin system but not from glutaredoxin and play an important role in eliminating peroxides generated during metabolism. PRDX1 might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H2O2. It reduces an intramolecular disulfide bond in GDPD5 that gates the ability to GDPD5 to drive postmitotic motor neuron differentiation. It may contribute to the antiviral activity of CD8(+) T-cells and have a proliferative effect in cancer development or progression.

本产品仅供科研使用，不用于临床诊断。