Recombinant Mouse B7-H4 (C-Fc)

货号：RB91EPT204

gui_ge

10ug

biao_da_su_zhu

Human Cells

biao_qian

C-Fc

fen_zi_liang

52.2 KDa

biao_guan_fen_zi_liang

70-90 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

huo_xing

Immobilized Anti-Human B7-H4 mAb(Cat#NC054) at 2μg/ml (100 μl/well) can bind Mouse B7-H4-Fc*(Cat#CS04).*: Biotinylated by NHS-biotin prior to testing The ED50 of Mouse B7-H4-Fc*(Cat#CS04) is 4.11 ng/ml.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Mouse V-set domain-containing T-cell activation inhibitor 1/VTCN1/B7-H4 is glycosylated member of the B7 family of immune co-stimulatory proteins. B7-H4 consists of extracellular domain (ECD) with one Ig-like V-set domain and one Ig-like C2-set domain. It is widely expressed, including in kidney, liver, lung, pancreas, placenta, prostate, spleen, testis and thymus. B7-H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. When expressed on the cell surface of tumor macrophages, plays an important role, together with regulatory T-cells (Treg), in the suppression of tumor-associated antigen-specific T-cell immunity. It also involved in promoting epithelial cell transformation.

本产品仅供科研使用，不用于临床诊断。