Recombinant Human CCL5

货号: RB91EPT210

gui_ge

10ug

biao_da_su_zhu

E.coli

biao_qian

NA

fen_zi_liang

7.8 KDa

biao_guan_fen_zi_liang

9-12 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Human Chemokine (C-C Motif) Ligand 5 (CCL5) plays an active role in recruiting leukocytes into inflammatory sites. CCL5 is secreted by many cell types at inflammatory sites and it exerts a wide range of activities through the receptors CCR1, CCR3, CCR4, and CCR5. N-Terminal truncated CCL5/RANTES, Met-RANTES, and amino-oxypentane (AOP)-RANTES exhibit antagonist or partial agonist functions on their receptors. CCL5/RANTES attracts different subtypes of leukocytes into inflamed tissue and intervenes in a wide range of allergic and autoimmune diseases.

本产品仅供科研使用，不用于临床诊断。