Recombinant Mouse Thrombomodulin (C-6His)

货号: RB91EPT230

gui_ge

10ug

biao_da_su_zhu

Human Cells

biao_qian

C-6His

fen_zi_liang

54.6 KDa

biao_guan_fen_zi_liang

85-90 KDa, reducing conditions

xing_tai

Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

chu_cun_fang_fa

Store at ≤-70° C, stable for 6 months after receipt.
Store at ≤-70° C, stable for 3 months under sterile conditions after opening.
Please minimize freeze-thaw cycles.

yun_shu_tiao_jian

The product is shipped on dry ice/polar packs.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Thrombomodulin is also known as CD141 antigen and blood dendritic cell antigen 3 (BDCA3), which is encoded by the THBD gene. The deduced amino acid sequence of mouse THBD predicts a signal peptide (aa 1 to 16) and a mature chain (aa 17 to 577) that consists of the following domains: C-type lectin, EGF-like, transmembrane and cytoplasmic. Mouse THBD is corresponding to the extracellular portion of the type I membrane protein. Predominantly synthesized by vascular endothelial cells, THBD inhibits coagulation and fibrinolysis. It functions as a cell surface receptor and an essential cofactor for active thrombin, which in turn activates protein C and thrombinactivatable fibrinolysis inhibitor (TAFI), also known as carboxypeptidase B2 (CPB2). In addition, THBD gene polymorphisims are associated with human disease and THBD plays a role in thrombosis, stroke, arteriosclerosis, and cancer.

本产品仅供科研使用，不用于临床诊断。