Recombinant Rat GM-CSF (C-6His)

货号: RB91EPT254

gui_ge

10ug

biao_da_su_zhu

Human Cells

biao_qian

C-6His

fen_zi_liang

15.6 KDa

biao_guan_fen_zi_liang

17-25 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF) was initially characterized as a growth factorthat can support the in vitro colony formation of granulocyte-macrophage progenitors. It is produced by anumber of different cell types (including activated T cells, B cells, macrophages, mast cells, endothelial cellsand fibroblasts) in response to cytokine of immune and inflammatory stimuli. Besides granulocyte-macrophageprogenitors, GM-CSF is also a growth factor for erythroid, megakaryocyte and eosinophil progenitors. Onmature hematopoietic, monocytes/ macrophages and eosinophils. GM-CSF has a functional role on nonhematopoitic cells. It can induce human endothelial cells to migrate and proliferate. Additionally, GM-CSF canalso stimulate the proliferation of a number of tumor cell lines, including osteogenic sarcoma, carcinoma andadenocarcinoma cell lines.

本产品仅供科研使用，不用于临床诊断。