Recombinant Human ACE-2 (C-mFc)

货号: RB91EPT272

gui_ge

50ug

biao_da_su_zhu

Human Cells

biao_qian

C-mFc

fen_zi_liang

110 KDa

biao_guan_fen_zi_liang

110-140 KDa, reducing conditions

xing_tai

Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 300mM NaCl, 1mM ZnCl₂, 10% Glycerol, pH 7.4.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

huo_xing

Immobilized 2019-nCoV S1 Protein (Cat#DRA35)at 10μg/ml (100μl/well) can bind Human ACE-2-mFc(Cat#C06A). The ED50 of Human ACE-2-mFc(Cat#C06A) is 45.11 ng/ml.

chu_cun_fang_fa

Store at ≤-70° C, stable for 6 months after receipt.
Store at ≤-70° C, stable for 3 months under sterile conditions after opening.
Please minimize freeze-thaw cycles.

yun_shu_tiao_jian

The product is shipped on dry ice/polar packs.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

Angiotensin-Converting Enzyme 2 (ACE-2) is an integral membrane protein and a zinc metalloprotease of the ACE family, the ACE family includes somatic and germinal ACE. ACE-2 cleaves angiotensins I and II as a carboxypeptidase, ACE-2 converts angiotensin I to angiotensin 1-9, and angiotensin II to angiotensin 1-7. ACE-2 is also able to hydrolyze apelin-13 and dynorphin-13 with high efficiency. ACE-2 can be high expressed in testis, kidney and heart, in colon, small intestine and ovary at moderate levels. Captopril and lisinopril as the classical ACE inhibitor don’t inhibit ACE-2 activity. ACE-2 may play an important role in regulating the heart function.

本产品仅供科研使用，不用于临床诊断。