Recombinant Human CD5L (C-6His)

货号: RB91EPT306

gui_ge

10ug

biao_da_su_zhu

Human Cells

biao_qian

C-6His

fen_zi_liang

37.09 KDa

biao_guan_fen_zi_liang

38-50 KDa, reducing conditions

xing_tai

Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

chun_du

Greater than 95% as determined by reducing SDS-PAGE.

nei_du_su_shui_ping

Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.

rong_jie_fang_fa

Always centrifuge tubes before opening.Do not mix by vortex or pipetting.
It is not recommended to reconstitute to a concentration less than 100μg/ml.
Dissolve the lyophilized protein in distilled water.
Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

chu_cun_fang_fa

Lyophilized protein should be stored at < -20° C, though stable at room temperature for 3 weeks.
Reconstituted protein solution can be stored at 4-7° C for 2-7 days.
Aliquots of reconstituted samples are stable at < -20° C for 3 months.

yun_shu_tiao_jian

The product is shipped at ambient temperature.
Upon receipt, store it immediately at the temperature listed below.

bei_jing_shuo_ming

CD5 Antigen-Like (CD5L) is a soluble protein that belongs to group B of the scavenger receptor cysteine-rich (SRCR) superfamily and contains three SRCR domains. CD5L is a secreted glycoprotein and expressed by macrophages presentin lymphoid tissues. It binds to myelomonocytic and lymphoid cells and may play an important role in the regulation of the innate and adaptive immune systems. CD5L functions as a pattern recognition molecule by binding both lipoteichoic acid (LTA) on Gram positive and lipopolysaccharide (LPS) on Gram-negative bacteria and the SRCR domain one of CD5L retains both the LPS and LTA binding activities. Furthermore, CD5L seems to play a role as an inhibitor of apoptosis.

本产品仅供科研使用，不用于临床诊断。