抗活化Caspase-3抗体

Anti-Cleaved Caspase-3 antibody - BSA and Azide free [Rabbit]

货号: RB9A37014879

产品描述

中文名称
重组Anti-Cleaved Caspase-3抗体 - BSA and Azide free(RB9A37014879)

英文名称
Anti-Cleaved Caspase-3 antibody - BSA and Azide free [Rabbit]

别名
CPP32 CASP3 Caspase-3 CASP-3 Apopain Cysteine protease CPP32 Protein Yama SREBP cleavage activity 1 CPP-32 SCA-1

靶点符号
CASP3

产品类型
一抗

关键信息

宿主种属
Rabbit

克隆
Monoclonal

亚型
IgG

不含载体蛋白
Yes

反应种属
Human

应用
WB ICC/IF

免疫原
The exact immunogen used to generate this antibody is proprietary information.

特异性
This antibody is more sensitive for the detection of cleaved caspase-3 than for pro-caspase-3. For better observation of pro-caspase-3, we recommend RB9A37051581.PubMedID 19789217 describes the detection of human cells injected into mice. The pro-caspase-3 is cleaved only when apoptosis event occurs. So, in order to detect active Caspase-3, we strongly suggest to induce your samples into apoptotic pathway.

反应性数据

应用	应用代码	物种	验证状态	备注
Flow Cytometry	FlowCyt	Human	不推荐	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Human	已验证	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	不推荐	-
Immunoprecipitation	IP	Human	不推荐	-
Western blot	WB	Human	已验证	The pro-caspase-3 is cleaved only when apoptosis event occurs. So, in order to detect active Caspase-3, we strongly suggest to induce your samples into apoptotic pathway.
Flow Cytometry	FlowCyt	Recombinant fragment	不推荐	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Recombinant fragment	不推荐	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Recombinant fragment	不推荐	-
Immunoprecipitation	IP	Recombinant fragment	不推荐	-
Western blot	WB	Recombinant fragment	不推荐	-
Flow Cytometry	FlowCyt	Recombinant fragment - Human	不推荐	-

应用	应用代码	物种	验证状态	备注
Immunocytochemistry/ Immunofluorescence	ICCIF	Recombinant fragment - Human	不推荐	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Recombinant fragment - Human	不推荐	-
Immunoprecipitation	IP	Recombinant fragment - Human	不推荐	-
Western blot	WB	Recombinant fragment - Human	已验证	The pro-caspase-3 is cleaved only when apoptosis event occurs. So, in order to detect active Caspase-3, we strongly suggest to induce your samples into apoptotic pathway.

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein A

存储溶液
pH: 7.2 - 7.4 Constituents: PBS

储存信息
Do Not Freeze

本产品仅供科研使用，不用于临床诊断。