# 核仁素抗体

Anti-Nucleolin antibody - BSA and Azide free [IgG1]

货号：RB9A37015784

## 产品描述

中文名称
Anti-Nucleolin抗体 - BSA and Azide free(RB9A37015784)

英文名称
Anti-Nucleolin antibody - BSA and Azide free [IgG1]

别名
Nucleolin　Protein C23　NCL

靶点符号
NCL

产品类型
一抗

## 关键信息

宿主种属
Mouse

克隆
Monoclonal

亚型
IgG1

轻链类型
kappa

不含载体蛋白
Yes

反应种属
Human

应用
IHC-P

免疫原
Cell preparation containing NCL protein. The exact immunogen used to generate this antibody is proprietary information.; P19338

浓度
1 mg/mL

## 反应性数据

| 应用 | 应用代码 | 物种 | 验证状态 | 推荐稀释 | 备注 |
| --- | --- | --- | --- | --- | --- |
| Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) | IHCP | Cow | 不推荐 | 0.25-0.5 µg/µL | (Primary incubation for 30 minutes at RT). Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. |
| Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) | IHCP | Human | 已验证 | 0.25-0.5 µg/mL | (Primary incubation for 30 minutes at RT). Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. |
| Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) | IHCP | Mouse | 不推荐 | 0.25-0.5 µg/mL | (Primary incubation for 30 minutes at RT). Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. |
| Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) | IHCP | Rat | 不推荐 | 0.25-0.5 µg/mL | (Primary incubation for 30 minutes at RT). Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. |

## 储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein G

纯化说明
RB9A37015784 was purified from Bioreactor Concentrate by Protein A/G.

存储溶液
pH: 7.2 - 7.4 Constituents: PBS

| 推荐的短期储存时间 | 储存信息 |
| --- | --- |
| 1-2 weeks | Avoid freeze / thaw cycle |

本产品仅供科研使用，不用于临床诊断。