抗p75神经营养因子受体抗体

Anti-p75 NGF Receptor antibody - BSA and Azide free [IgG1]

货号: RB9A37015786

产品描述

中文名称

Anti-p75 NGF Receptor抗体 - BSA and Azide free(RB9A37015786)

英文名称

Anti-p75 NGF Receptor antibody - BSA and Azide free [IgG1]

别名

CD271 TNFRSF16 NGFR Tumor necrosis factor receptor superfamily member 16 Gp80-LNGFR Low affinity neurotrophin receptor p75NTR Low-affinity nerve growth factor receptor Low-affinity nerve growth factor receptor p75NGFR Low-affinity nerve growth factor receptor p75NGR p75 ICD NGF receptor

靶点符号

NGFR

产品类型

一抗

关键信息

宿主种属

Mouse

克隆

Monoclonal

亚型

IgG1

轻链类型

kappa

不含载体蛋白

Yes

反应种属

Human

应用

IHC-P

免疫原

Cell preparation containing NGFR protein. The exact immunogen used to generate this antibody is proprietary information.; P08138

浓度

1 mg/mL

表位

Epitope spans aa 1-160 of the extracellular domain of NGFR/NTR.

反应性数据

应用	应用代码	物种	验证状态	推荐稀释	备注
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	已验证	0.5-1 µg/mL	for 30 minutes at room temperature. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Mouse	不推荐	0.5-1 µg/mL	for 30 minutes at room temperature. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rat	不推荐	0.5-1 µg/mL	for 30 minutes at room temperature. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein G

纯化说明
Purified from bioreactor concentrate.

存储溶液
pH: 7.2 - 7.4 Constituents: PBS

推荐的短期储存时间
1-2 weeks

储存信息
Avoid freeze / thaw cycle

本产品仅供科研使用，不用于临床诊断。