抗Bcl-2抗体

Anti-Bcl-2 antibody - BSA and Azide free [Mouse]

货号：RB9A37015814

产品描述

中文名称
Anti-Bcl-2抗体 - BSA and Azide free

英文名称
Anti-Bcl-2 antibody - BSA and Azide free [Mouse]

别名
Apoptosis regulator Bcl-2 BCL2

靶点符号
BCL2

产品类型
一抗

关键信息

宿主种属
Mouse

克隆
Monoclonal

亚型
IgG1

轻链类型
kappa

不含载体蛋白
Yes

反应种属
Human Cat Horse Pig Llama

应用
Flow Cyt WB IHC-P

免疫原
Synthetic Peptide within Human BCL2 aa 1-100. The exact immunogen used to generate this antibody is proprietary information.; P10415

浓度
1 mg/mL

反应性数据

应用	应用代码	物种	验证状态	备注	推荐稀释
Flow Cytometry	FlowCyt	Cat	保证反应	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Cat	已验证	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.	-
Western blot	WB	Cat	保证反应	-	-
Flow Cytometry	FlowCyt	Cow	不推荐	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Cow	不推荐	-	-
Western blot	WB	Cow	不推荐	-	-
Flow Cytometry	FlowCyt	Horse	保证反应	-	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Horse	已验证	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.	-
Western blot	WB	Horse	保证反应	-	-
Flow Cytometry	FlowCyt	Human	已验证	(in 0.1ml volume)	1-2 µg for 10^6 Cells
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	已验证	(for 30 minutes at RT) Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.	1-2 µg/mL
Western blot	WB	Human	已验证	-	1-2 µg/mL
Flow Cytometry	FlowCyt	Llama	保证反应	-	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Llama	已验证	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.	-
Western blot	WB	Llama	保证反应	-	-
Flow Cytometry	FlowCyt	Mouse	不推荐	(in 0.1ml volume)	1-2 µg for 10^6 Cells
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Mouse	不推荐	(for 30 minutes at RT) Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.	1-2 µg/mL
Western blot	WB	Mouse	不推荐	-	1-2 µg/mL
Flow Cytometry	FlowCyt	Pig	保证反应	-	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Pig	已验证	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.	-
Western blot	WB	Pig	保证反应	-	-
Flow Cytometry	FlowCyt	Rat	不推荐	(in 0.1ml volume)	1-2 µg for 10^6 Cells
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rat	不推荐	(for 30 minutes at RT) Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.	1-2 µg/mL
Western blot	WB	Rat	不推荐	-	1-2 µg/mL

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein A/G

纯化说明
RB9A37015814 is purified from Bioreactor Concentrate by Protein A/G.

存储溶液
pH: 7.2 - 7.4 Constituents: PBS

推荐的短期储存时间	储存信息
1-2 weeks	Avoid freeze / thaw cycle

本产品仅供科研使用，不用于临床诊断。