波形蛋白抗体

Anti-Vimentin antibody - BSA and Azide free [IgG1]

货号：RB9A37015837

产品描述

中文名称
Anti-Vimentin抗体 - BSA and Azide free(RB9A37015837)

英文名称
Anti-Vimentin antibody - BSA and Azide free [IgG1]

别名
Vimentin VIM

靶点符号
VIM

产品类型
一抗

关键信息

宿主种属
Mouse

克隆
Monoclonal

亚型
IgG1

轻链类型
kappa

不含载体蛋白
Yes

反应种属
Human

应用
IHC-P WB Protein Array

免疫原
Recombinant Full Length Protein corresponding to Human VIM.; P08670

特异性
This antibody shows no cross-reaction with other closely related intermediate filament proteins (IFP s) such as desmin, keratin, neurofilament, and glial fibrillary acid protein.

浓度
1 mg/mL

反应性数据

应用	应用代码	物种	验证状态	推荐稀释	备注
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	已验证	1-2 μg/mL	Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes. Incubate with primary antibody for 30 mins at RT.
Protein Array	ProteinArray	Human	已验证	-	-
Western blot	WB	Human	已验证	1-22 μg/mL	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Mouse	不推荐	1-2 μg/mL	Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes. Incubate with primary antibody for 30 mins at RT.
Protein Array	ProteinArray	Mouse	不推荐	-	-
Western blot	WB	Mouse	不推荐	1-2 μg/mL	-

应用	应用代码	物种	验证状态	推荐稀释	备注
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rat	不推荐	1-2 μg/mL	Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes. Incubate with primary antibody for 30 mins at RT.
Protein Array	ProteinArray	Rat	不推荐	-	-
Western blot	WB	Rat	不推荐	1-2 μg/mL	-

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein A/G

纯化说明
Purified from bioreactor concentrate.

存储溶液
pH: 7.2 - 7.4 Constituents: PBS

推荐的短期储存时间
1-2 weeks

储存信息
Avoid freeze / thaw cycle

本产品仅供科研使用，不用于临床诊断。