抗核因子κB受体活化因子配体抗体

Anti-RANKL antibody (RB9A37016572

货号：RB9A37016572

产品描述

中文名称
Anti-RANKL抗体 (RB9A37016572)

英文名称
Anti-RANKL antibody (RB9A37016572)

别名
CD254OPGLRANKLTRANCETNFSF11Tumor necrosis factor ligand superfamily member 11Osteoclast differentiation factorOsteoprotegerin ligandReceptor activator of nuclear factor kappa-B ligandTNF-related activation-induced cytokineODF

靶点符号
TNFSF11

产品类型
一抗

关键信息

宿主种属
Rabbit

克隆
Polyclonal

亚型
IgG

不含载体蛋白
No

反应种属
MouseHuman

应用
IHC-PFlow Cyt (Intra)

免疫原
Synthetic Peptide within Human TNFSF11 aa 250-350 conjugated to Keyhole Limpet Haemocyanin. The exact immunogen used to generate this antibody is proprietary information.; O14788

浓度
1 mg/mL

反应性数据

应用	应用代码	物种	验证状态	推荐稀释	备注
Flow Cytometry (Intracellular)	FlowCytIntra	Human	已验证	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	已验证	1/100 - 1/500	Boil slides in 0.01M sodium citrate buffer (pH6) at 100° C for 15-20 minutes. Remove the slides from heat and allow them to stand at RT in buffer for 20 minutes. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Flow Cytometry (Intracellular)	FlowCytIntra	Mouse	保证反应	-	-

应用	应用代码	物种	验证状态	推荐稀释	备注
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Mouse	已验证	1/100 - 1/500	Boil slides in 0.01M sodium citrate buffer (pH6) at 100° C for 15-20 minutes. Remove the slides from heat and allow them to stand at RT in buffer for 20 minutes. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein A

存储溶液
pH: 7.4 Preservative: 0.02% Proclin 300 Constituents: 50% Glycerol (glycerin, glycerine), 48.98% TBS, 1X, 1% BSA

推荐的短期储存时间
1-2 weeks

储存信息
Avoid freeze / thaw cycle

本产品仅供科研使用，不用于临床诊断。