上皮细胞黏附分子抗体

Anti-EpCAM/CD326 antibody [Rabbit]

货号：RB9A37016675

产品描述

中文名称
Anti-EpCAM/CD326抗体（RB9A37016675）

英文名称
Anti-EpCAM/CD326 antibody [Rabbit]

别名
CD326Tacstd1EpcamEpithelial cell adhesion moleculeEp-CAMEpithelial glycoprotein 314Protein 289ATumor-associated calcium signal transducer 1EGP314mEGP314

靶点符号
EpCAM

产品类型
一抗

关键信息

宿主种属
Rabbit

克隆
Polyclonal

亚型
IgG

不含载体蛋白
No

反应种属
MouseRatRabbit

应用
WBIHC-P

免疫原
Synthetic Peptide within Mouse EpCAM/CD326 aa 50-150 conjugated to Keyhole Limpet Haemocyanin. The exact immunogen used to generate this antibody is proprietary information.; Q99JW5

浓度
1 mg/mL

反应性数据

应用	应用代码	物种	验证状态	推荐稀释	备注
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	预测反应	-	-
Western blot	WB	Human	预测反应	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Mouse	保证反应	-	-
Western blot	WB	Mouse	已验证	1/100 - 1/1000	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rabbit	已验证	1/100 - 1/500	When using a fluorescent probe the recommended dilution is 1/50 - 1/200. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Western blot	WB	Rabbit	保证反应	-	-

应用	应用代码	物种	验证状态	推荐稀释	备注
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rat	已验证	1/100 - 1/500	When using a fluorescent probe the recommended dilution is 1/50 - 1/200. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Western blot	WB	Rat	保证反应	-	-

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein A

存储溶液
pH: 7.4 Preservative: 0.02% Proclin 300 Constituents: 50% Glycerol (glycerin, glycerine), 48.98% TBS, 1X, 1% BSA

推荐的短期储存时间
1-2 weeks

储存信息
Avoid freeze / thaw cycle

本产品仅供科研使用，不用于临床诊断。