抗Iba1抗体

Anti-Iba1 antibody - BSA and Azide free (RB9A37017538

货号：RB9A37017538

产品描述

中文名称
重组Anti-Iba1抗体 - BSA and Azide free(RB9A37017538)

英文名称
Anti-Iba1 antibody - BSA and Azide free (RB9A37017538)

别名
G1IBA1AIF1Allograft inflammatory factor 1AIF-1Ionized calcium-binding adapter molecule 1Protein G1iba1iba-1iba 1

靶点符号
AIF1

产品类型
一抗

关键信息

宿主种属
Rabbit

克隆
Monoclonal

亚型
IgG

不含载体蛋白
Yes

反应种属
MouseRatHuman

应用
ICC/IFIHC-FrIHC-PFlow Cyt (Intra)WB

免疫原
The exact immunogen used to generate this antibody is proprietary information.

特异性
For RB9A37008469 Abcam recommends blocking in 3% milk for cleanest results in WB. Blocking with BSA gives slightly higher background.

反应性数据

应用	应用代码	物种	验证状态	备注
Flow Cytometry (Intracellular)	FlowCytIntra	Human	已验证	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Human	保证反应	-
Immunohistochemistry (Frozen sections)	IHCFr	Human	保证反应	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	已验证	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
Western blot	WB	Human	已验证	Abcam recommends blocking in 3% milk for cleanest results in WB. Blocking with BSA gives slightly higher background.
Flow Cytometry (Intracellular)	FlowCytIntra	Mouse	已验证	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Mouse	已验证	-
Immunohistochemistry (Frozen sections)	IHCFr	Mouse	已验证	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Mouse	已验证	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

应用	应用代码	物种	验证状态	备注
Western blot	WB	Mouse	已验证	Abcam recommends blocking in 3% milk for cleanest results in WB. Blocking with BSA gives slightly higher background.
Flow Cytometry (Intracellular)	FlowCytIntra	Rat	保证反应	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Rat	保证反应	-
Immunohistochemistry (Frozen sections)	IHCFr	Rat	已验证	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rat	已验证	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
Western blot	WB	Rat	已验证	Abcam recommends blocking in 3% milk for cleanest results in WB. Blocking with BSA gives slightly higher background.

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein A

存储溶液
pH: 7.2 - 7.4 Constituents: PBS

储存信息
Do Not Freeze

本产品仅供科研使用，不用于临床诊断。