抗血小板内皮细胞黏附分子1抗体

Anti-PECAM-1/CD31 antibody - BSA and Azide free (RB9A37019070

货号: RB9A37019070

产品描述

中文名称
重组Anti-PECAM-1/CD31抗体 - BSA and Azide free(RB9A37019070)

英文名称
Anti-PECAM-1/CD31 antibody - BSA and Azide free (RB9A37019070)

别名
CD31Platelet endothelial cell adhesion moleculePECAM-1EndoCAMGPIIA'PECA1PECAM1

靶点符号
PECAM-1

产品类型
一抗

关键信息

宿主种属
Rabbit

克隆
Monoclonal

亚型
IgG

不含载体蛋白
Yes

反应种属
MouseRatHuman

应用
mIHCIHC-PIHC-FoFr

免疫原
The exact immunogen used to generate this antibody is proprietary information.

反应性数据

应用	应用代码	物种	验证状态	备注
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Human	保证反应	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	已验证	In our hands we observed non-specific cytoplasmic staining on tubular cells in rat kidney. The ideal fixation time will depend on the size of the tissue block and the type of tissue, but fixation between 18 - 24h is suitable for most samples. Positive Control: human lung tissue. Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
Multiplex immunohistochemistry	mIHC	Human	已验证	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Mouse	已验证	-

应用	应用代码	物种	验证状态	备注
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Mouse	已验证	In our hands we observed non-specific cytoplasmic staining on tubular cells in rat kidney. The ideal fixation time will depend on the size of the tissue block and the type of tissue, but fixation between 18 - 24h is suitable for most samples. Positive Control: human lung tissue. Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
Multiplex immunohistochemistry	mIHC	Mouse	已验证	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Rat	已验证	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rat	已验证	In our hands we observed non-specific cytoplasmic staining on tubular cells in rat kidney. The ideal fixation time will depend on the size of the tissue block and the type of tissue, but fixation between 18 - 24h is suitable for most samples. Positive Control: human lung tissue. Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

应用	应用代码	物种	验证状态	备注
Multiplex immunohistochemistry	mIHC	Rat	已验证	-

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Protein A

存储溶液
pH: 7.2 - 7.4 Constituents: PBS

储存信息
Do Not Freeze

本产品仅供科研使用，不用于临床诊断。