血管性血友病因子抗体

Anti-Von Willebrand Factor antibody (RB9A37057196

货号：RB9A37057196

产品描述

中文名称

Anti-Von Willebrand Factor 抗体（RB9A37057196）

英文名称

Anti-Von Willebrand Factor antibody (RB9A37057196)

别名

F8VWF VWF von Willebrand factor vWF

靶点符号

VWF

产品类型

一抗

关键信息

宿主种属

Rabbit

克隆

Polyclonal

亚型

IgG

不含载体蛋白

No

反应种属

Human Sheep Rat Pig Horse Dog Guinea pig Cow

应用

Flow Cyt IHC-FrFl IHC-FoFr ICC/IF WB IHC-Fr IHC-P

免疫原

Native Full Length Protein corresponding to Human VWF.; P04275

反应性数据

应用	应用代码	物种	验证状态	备注	推荐稀释
Flow Cytometry	FlowCyt	Chicken	不推荐	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunocytochemistry/ Immunofluorescence	ICCIF	Chicken	不推荐	-	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Chicken	不推荐	-	-
Immunohistochemistry (Frozen sections)	IHCFr	Chicken	不推荐	-	-
Immunohistochemistry - Free Floating	IHCFrFl	Chicken	不推荐	(see Abreview)	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Chicken	不推荐	for IF and 1/1000-1/2000 for ABC methods with HRP conjugates. Perform enzymatic antigen retrieval with 0.1% pronase for 10 min at 35 ° C before commencing with IHC protocol.Indirect Immunofluorescence: minimum working dilution of 1:200 was determined using FFPE sections of human tongue with FITC-conjugated secondary.Indirect Immunoperoxidase Labeling: minimum working dilution of 1:800 was determined using FFPE sections of human tongue with biotinylated secondary and signal amplification.	-
Western blot	WB	Chicken	不推荐	-	-
Flow Cytometry	FlowCyt	Cow	保证反应	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunocytochemistry/ Immunofluorescence	ICCIF	Cow	保证反应	-	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Cow	保证反应	-	-
Immunohistochemistry (Frozen sections)	IHCFr	Cow	保证反应	-	-
Immunohistochemistry - Free Floating	IHCFrFl	Cow	保证反应	(see Abreview)	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Cow	不推荐	-	-
Western blot	WB	Cow	保证反应	-	-
Flow Cytometry	FlowCyt	Dog	保证反应	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Dog	保证反应	-	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Dog	保证反应	-	-
Immunohistochemistry (Frozen sections)	IHCFr	Dog	保证反应	-	-
Immunohistochemistry - Free Floating	IHCFrFl	Dog	保证反应	(see Abreview)	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Dog	不推荐	-	-
Western blot	WB	Dog	保证反应	-	-
Flow Cytometry	FlowCyt	Guinea pig	保证反应	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Guinea pig	保证反应	-	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Guinea pig	保证反应	-	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (Frozen sections)	IHCFr	Guinea pig	保证反应	-	-
Immunohistochemistry - Free Floating	IHCFrFl	Guinea pig	保证反应	(see Abreview)	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Guinea pig	不推荐	-	-
Western blot	WB	Guinea pig	保证反应	-	-
Flow Cytometry	FlowCyt	Horse	保证反应	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Horse	保证反应	-	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Horse	保证反应	-	-
Immunohistochemistry (Frozen sections)	IHCFr	Horse	保证反应	-	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry - Free Floating	IHCFrFl	Horse	保证反应	(see Abreview)	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Horse	不推荐	-	-
Western blot	WB	Horse	保证反应	-	-
Flow Cytometry	FlowCyt	Human	保证反应	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Human	保证反应	-	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Human	保证反应	-	-
Immunohistochemistry (Frozen sections)	IHCFr	Human	保证反应	-	-
Immunohistochemistry - Free Floating	IHCFrFl	Human	保证反应	(see Abreview)	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	已验证	for IF and 1/1000-1/2000 for ABC methods with HRP conjugates. Perform enzymatic antigen retrieval with 0.1% pronase for 10 min at 35 ° C before commencing with IHC protocol.Indirect Immunofluorescence: minimum working dilution of 1:200 was determined using FFPE sections of human tongue with FITC-conjugated secondary.Indirect Immunoperoxidase Labeling: minimum working dilution of 1:800 was determined using FFPE sections of human tongue with biotinylated secondary and signal amplification.	1/200 - 1/400
Western blot	WB	Human	保证反应	-	-
Flow Cytometry	FlowCyt	Pig	保证反应	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunocytochemistry/ Immunofluorescence	ICCIF	Pig	保证反应	-	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Pig	保证反应	-	-
Immunohistochemistry (Frozen sections)	IHCFr	Pig	保证反应	-	-
Immunohistochemistry - Free Floating	IHCFrFl	Pig	保证反应	(see Abreview)	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Pig	不推荐	-	-
Western blot	WB	Pig	保证反应	-	-
Flow Cytometry	FlowCyt	Rat	保证反应	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Rat	保证反应	-	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Rat	保证反应	-	-
Immunohistochemistry (Frozen sections)	IHCFr	Rat	保证反应	-	-
Immunohistochemistry - Free Floating	IHCFrFl	Rat	保证反应	(see Abreview)	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rat	不推荐	-	-
Western blot	WB	Rat	保证反应	-	-
Flow Cytometry	FlowCyt	Sheep	保证反应	RB9A37007601 - Rabbit polyclonal IgG, is suitable for use as an isotype control with this antibody.	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Sheep	保证反应	-	-
Immunohistochemistry (PFA perfusion fixed frozen sections)	IHCFoFr	Sheep	保证反应	-	-

应用	应用代码	物种	验证状态	备注	推荐稀释
Immunohistochemistry (Frozen sections)	IHCFr	Sheep	保证反应	-	-
Immunohistochemistry - Free Floating	IHCFrFl	Sheep	保证反应	(see Abreview)	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Sheep	不推荐	-	-
Western blot	WB	Sheep	保证反应	-	-

储存与纯化

形式
Liquid

纯化说明
Whole antiserum is fractionated and then further purified by ion exchange chromatography to provide the IgG fraction of antiserum. This fraction is essentially free of other rabbit serum proteins.

存储溶液
pH: 7.4 Preservative: 0.097% Sodium azide Constituents: PBS

推荐的短期储存时间
1-2 weeks

储存信息
Avoid freeze / thaw cycle

纯度
IgG fraction

本产品仅供科研使用，不用于临床诊断。