抗黑色素瘤gp100抗体

Anti-Melanoma gp100 antibody (RB9A37058357

货号：RB9A37058357

产品描述

中文名称

Anti-Melanoma gp100 抗体（RB9A37058357）

英文名称

Anti-Melanoma gp100 antibody (RB9A37058357)

别名

D12S53E PMEL17 SILV PMEL Melanocyte protein PMEL ME20-M Melanocyte protein Pmel 17 Melanocytes lineage-specific antigen GP100 Melanoma-associated ME20 antigen P1 P100 Premelanosome protein Silver locus protein homolog ME20M

靶点符号

PMEL

产品类型

一抗

关键信息

宿主种属

Mouse

克隆

Monoclonal

亚型

IgG1

轻链类型

kappa

不含载体蛋白

No

反应种属

Human

应用

IHC-P Flow Cyt ICC/IF

免疫原

Cell preparation containing PMEL protein. The exact immunogen used to generate this antibody is proprietary information.; P40967

反应性数据

应用	应用代码	物种	验证状态	推荐稀释	备注
Flow Cytometry	FlowCyt	Dog	不推荐	1/10	RB9A37007391 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
Immunocytochemistry/ Immunofluorescence	ICCIF	Dog	不推荐	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Dog	不推荐	0.5-1 µg/mL	Incubate for 30 minutes at room temperature. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes
Flow Cytometry	FlowCyt	Human	已验证	1/10	RB9A37007391 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
Immunocytochemistry/ Immunofluorescence	ICCIF	Human	保证反应	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	已验证	0.5-1 µg/mL	Incubate for 30 minutes at room temperature. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes

应用	应用代码	物种	验证状态	推荐稀释	备注
Flow Cytometry	FlowCyt	Rat	不推荐	1/10	RB9A37007391 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
Immunocytochemistry/ Immunofluorescence	ICCIF	Rat	不推荐	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Rat	不推荐	0.5-1 µg/mL	Incubate for 30 minutes at room temperature. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes

储存与纯化

形式
Liquid

存储溶液
pH: 7.2 - 7.4 Preservative: 0.05% Sodium azide Constituents: Tissue culture supernatant, 0.05% BSA

推荐的短期储存时间
1-2 weeks

储存信息
Avoid freeze / thaw cycle

纯度
Tissue culture supernatant

本产品仅供科研使用，不用于临床诊断。