抗核因子κB受体活化因子配体抗体

Anti-RANKL antibody [Rabbit]

货号：RB9A37060588

产品描述

中文名称
Anti-RANKL 抗体（RB9A37060588）

英文名称
Anti-RANKL antibody [Rabbit]

别名
CD254 OPGL RANKL TRANCE TNFSF11 Tumor necrosis factor ligand superfamily member 11 Osteoclast differentiation factor Osteoprotegerin ligand Receptor activator of nuclear factor kappa-B ligand TNF-related activation-induced cytokine ODF

靶点符号
TNFSF11

产品类型
一抗

关键信息

宿主种属
Rabbit

克隆
Polyclonal

亚型
IgG

不含载体蛋白
No

反应种属
Human

应用
ICC/IF ELISA IHC-P WB sELISA

免疫原
Recombinant Full Length Protein corresponding to Human TNFSF11.; O14788

反应性数据

应用	应用代码	物种	验证状态	推荐稀释	备注

应用	应用代码	物种	验证状态	推荐稀释	备注
ELISA	ELISA	Human	预测反应	-	-
Immunocytochemistry/ Immunofluorescence	ICCIF	Human	保证反应	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Human	保证反应	-	-
Western blot	WB	Human	已验证	1 μg/mL	To detect hsRANK-L by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hsRANK-L is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.
Sandwich ELISA	sELISA	Human	保证反应	-	-

应用	应用代码	物种	验证状态	推荐稀释	备注
ELISA	ELISA	Recombinant full length protein	保证反应	-	To detect hsRANK-L by direct ELISA (using 100µl/well antibody solution) a concentration of at least 0.5µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of 0.2 - 0.4 ng/well of recombinant hsRANK-L.
Immunocytochemistry/ Immunofluorescence	ICCIF	Recombinant full length protein	不推荐	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Recombinant full length protein	不推荐	-	-
Western blot	WB	Recombinant full length protein	不推荐	-	-
Sandwich ELISA	sELISA	Recombinant full length protein	不推荐	-	-
ELISA	ELISA	Recombinant full length protein - Human	不推荐	-	-

应用	应用代码	物种	验证状态	推荐稀释	备注
Immunocytochemistry/ Immunofluorescence	ICCIF	Recombinant full length protein - Human	不推荐	-	-
Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections)	IHCP	Recombinant full length protein - Human	不推荐	-	-
Western blot	WB	Recombinant full length protein - Human	不推荐	-	-
Sandwich ELISA	sELISA	Recombinant full length protein - Human	已验证	-	To detect hsRANKL by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with RB9A37058726 as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hsRANKL.

储存与纯化

形式

Lyophilized

纯化工艺

Affinity purification Immunogen

推荐的短期储存时间

1-2 weeks

复溶

Reconstitute with 200µl of sterile water. Please note that if you receive this product in liquid form it has already been reconstituted as described and no further reconstitution is necessary.

本产品仅供科研使用，不用于临床诊断。