Goat F

ab’

货号：RB9A37061809

产品描述

中文名称
山羊F(ab’)2抗人IgG - (Fab’)2 (DyLight® 488),预吸附二抗

英文名称
Goat F(ab’)2 Anti-Human IgG - (Fab’)2 (DyLight® 488), pre-adsorbed

别名
Immunoglobulin heavy constant gamma 1Ig gamma-1 chain C regionIg gamma-1 chain C region EUIg gamma-1 chain C region KOLIg gamma-1 chain C region NIEIGHG1Immunoglobulin heavy constant gamma 4Ig gamma-4 chain C regionIGHG4Immunoglobulin heavy constant gamma 3HDCHeavy chain disease proteinIg gamma-3 chain C regionIGHG3Immunoglobulin heavy constant gamma 2Ig gamma-2 chain C regionIg gamma-2 chain C region DOTIg gamma-2 chain C region TILIg gamma-2 chain C region ZIEIGHG2

靶点符号
IGHG1

产品类型
二抗

关键信息

宿主种属
Goat

克隆
Polyclonal

亚型
IgG

偶联物
DyLight® 488

激发波长/发射波长
Ex: 493nm, Em: 518nm

反应种属
Human

应用
ICC/IFFlow CytIHC-P

靶标种属
Human

靶标亚型
IgG

靶标特异性
F(ab')2

预吸附
Yes

最低交叉反应
Mouse, Rat,

浓度
0.5 mg/mL

反应性数据

应用	验证状态	推荐稀释
Flow Cyt	已验证	1/50 - 1/200
ICC/IF	已验证	1/50 - 1/500
IHC-P	已验证	1/50 - 1/500

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Immunogen

纯化说明
This antibody was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')2 fragment were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Fragments were conjugated to DyLight® 488.

存储溶液
pH: 6.8 - 7.4 Preservative: 0.09% Sodium azide Constituents: PBS, 0.2% BSA

本产品仅供科研使用，不用于临床诊断。