山羊抗人 IgG Fc

Goat Anti-Human IgG Fc (DyLight® 650

货号：RB9A37061820

产品描述

中文名称
Goat Anti-Human IgG Fc (DyLight® 650) preadsorbed

英文名称
Goat Anti-Human IgG Fc (DyLight® 650) preadsorbed

别名
Immunoglobulin heavy constant gamma 1Ig gamma-1 chain C regionIg gamma-1 chain C region EUIg gamma-1 chain C region KOLIg gamma-1 chain C region NIEIGHG1Immunoglobulin heavy constant gamma 4Ig gamma-4 chain C regionIGHG4Immunoglobulin heavy constant gamma 3HDCHeavy chain disease proteinIg gamma-3 chain C regionIGHG3Immunoglobulin heavy constant gamma 2Ig gamma-2 chain C regionIg gamma-2 chain C region DOTIg gamma-2 chain C region TILIg gamma-2 chain C region ZIEIGHG2

靶点符号
IGHG1

产品类型
二抗

关键信息

宿主种属
Goat

克隆
Polyclonal

亚型
IgG

偶联物
DyLight® 650

激发波长/发射波长
Ex: 654nm, Em: 673nm

反应种属
Human

应用
ICC/IFIHC-PFlow Cyt

靶标种属
Human

靶标亚型
IgG

靶标特异性
Fc region

预吸附
Yes

最低交叉反应
Mouse, Rat, Sheep, Rabbit, Goat, Horse, Chicken, Cow,

浓度
0.5 mg/mL

反应性数据

应用	验证状态	推荐稀释	备注
Flow Cyt	已验证	1/50 - 1/200	-
ICC/IF	已验证	1/50 - 1/500	-
IHC-P	已验证	1/50 - 1/500	DyLight ® 650 replaces DyLight ® 649. The DyLight ® 650 excitation is 652nm and the emission is 672nm. DyLight ® 650 hence provides the same far-red fluorescence and photostability as the DyLight ® 649 dye. Please refer to the vial in order to check whether the product you received is DyLight ® 649 or DyLight ® 650. This information however will have no impact on your experiments.

储存与纯化

形式
Liquid

纯化工艺
Affinity purification Immunogen

纯化说明	存储溶液
This antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 650.	pH: 6.8 - 7.4 Preservative: 0.09% Sodium azide Constituents: PBS, 0.2% BSA

本产品仅供科研使用，不用于临床诊断。