抗人LAP/TGF-β1参考抗体

Anti-Human LAP/TGF-beta 1 Reference Antibody

货号：RB9T37016163

在您使用本产品前请认真阅读说明书，这对您顺利完成实验很有帮助。

产品编码
RB9T37016163

中文名称
抗人LAP/TGF-β1参考抗体

英文名称
Anti-Human LAP/TGF-beta 1 Reference Antibody

形态
Liquid

纯度
>95% purity as determined by SDS-PAGE.

内毒素水平
Please contact the lab for this information.

应用
ELISA, Bioactivity: FACS, Functional assay, Research in vivo

表达系统
Mammalian Cells

登录号
P54831

宿主种属
Caninized

克隆性
Monoclonal

抗体亚型
IgG2, kappa

纯化方式
Protein A/G purified from cell culture supernatant.

浓度
1mg/ml

种属反应性
Dog

靶标

TGFB1/TGF-beta-1

背景信息

Cirevetmab (ZTS-00521426) is an immunoglobulin G2-kappa, Canis lupus familiaris TGFB1 caninized monoclonal antibody. Cirevetmab is an immunomodulator.

• TGFB1 induces fetal reprogramming and enhances intestinal regeneration., PMID:37865088

• Biomarkers for the identification of cardiac fibroblast and myofibroblast cells., PMID:29987445

• Association between TGF-β gene polymorphism and myopia: A systematic review and meta-analysis., PMID:35905284

• A fibroblast-dependent TGF-β1/sFRP2 noncanonical Wnt signaling axis promotes epithelial metaplasia in idiopathic pulmonary fibrosis., PMID:38980870

• The effect of TGF-β_1 polymorphisms on pulmonary disease progression in patients with cystic fibrosis., PMID:35525938

• Lack of association between the TGF-β(1) gene and development of COPD in Asians: a case-control study and meta-analysis., PMID:21556788

• TGF-β and Diabetic Nephropathy: Lessons Learned Over the Past 20 Years., PMID:32039767

• Direct effect of transforming growth factor-beta 1 (TGF-β1) on human apical papilla cell proliferation and mineralisation., PMID:34596309

• Concentration change of TGF -β 1 in aqueous humor of rabbits., PMID:24507648

• TGF-beta 1 codon 10 polymorphism is associated with cerebral SVD., PMID:22030425

别名

ZTS00521426, ZTS-00521426, ZTS00521426, 2473240-98-9

储存缓冲液

0.01M PBS, pH 7.4.

稳定性和储存

Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4° C for short-term storage (1-2 weeks). Store at -20° C for up to 12 months. For long-term storage, store at -80° C.

For research use only. Not suitable for clinical or therapeutic use.